

University of Groningen

## Cost-Effectiveness of Magnetic Resonance Angiography Versus Intra-arterial Digital Subtraction Angiography to Follow-Up Patients With Coiled Intracranial Aneurysms

Schaafsma, Joanna D.; Koffijberg, Hendrik; Buskens, Erik; Velthuis, Birgitta K.; van der Graaf, Yolanda; Rinkel, Gabriel J. E.

*Published in:*  
Stroke

*DOI:*  
[10.1161/STROKEAHA.110.585083](https://doi.org/10.1161/STROKEAHA.110.585083)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2010

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Schaafsma, J. D., Koffijberg, H., Buskens, E., Velthuis, B. K., van der Graaf, Y., & Rinkel, G. J. E. (2010). Cost-Effectiveness of Magnetic Resonance Angiography Versus Intra-arterial Digital Subtraction Angiography to Follow-Up Patients With Coiled Intracranial Aneurysms. *Stroke*, 41(8), 1736-1742. <https://doi.org/10.1161/STROKEAHA.110.585083>

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# Stroke

American Stroke  
Association<sup>SM</sup>

JOURNAL OF THE AMERICAN HEART ASSOCIATION

A Division of American  
Heart Association



## **Cost-Effectiveness of Magnetic Resonance Angiography Versus Intra-arterial Digital Subtraction Angiography to Follow-Up Patients With Coiled Intracranial Aneurysms**

Joanna D. Schaafsma, Hendrik Koffijberg, Erik Buskens, Birgitta K. Velthuis,  
Yolanda van der Graaf and Gabriël J.E. Rinkel

*Stroke* 2010;41;1736-1742; originally published online Jul 1, 2010;

DOI: 10.1161/STROKEAHA.110.585083

Stroke is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75214  
Copyright © 2010 American Heart Association. All rights reserved. Print ISSN: 0039-2499. Online  
ISSN: 1524-4628

The online version of this article, along with updated information and services, is  
located on the World Wide Web at:

<http://stroke.ahajournals.org/cgi/content/full/41/8/1736>

Subscriptions: Information about subscribing to Stroke is online at  
<http://stroke.ahajournals.org/subscriptions/>

Permissions: Permissions & Rights Desk, Lippincott Williams & Wilkins, a division of Wolters  
Kluwer Health, 351 West Camden Street, Baltimore, MD 21202-2436. Phone: 410-528-4050. Fax:  
410-528-8550. E-mail:  
[journalpermissions@lww.com](mailto:journalpermissions@lww.com)

Reprints: Information about reprints can be found online at  
<http://www.lww.com/reprints>

# Cost-Effectiveness of Magnetic Resonance Angiography Versus Intra-arterial Digital Subtraction Angiography to Follow-Up Patients With Coiled Intracranial Aneurysms

Joanna D. Schaafsma, MD; Hendrik Koffijberg, PhD; Erik Buskens, MD, PhD;  
Birgitta K. Velthuis, MD, PhD; Yolanda van der Graaf, MD, PhD; Gabriël J.E. Rinkel, MD

**Background and Purpose**—To follow up patients with coiled intracranial aneurysms, magnetic resonance angiography (MRA) is a promising noninvasive alternative to current standard intra-arterial digital subtraction angiography (IA-DSA). MRA test results do not always concord with those of IA-DSA, and the impact of discrepancies on health benefits and costs is unknown. We evaluated the cost-effectiveness of follow-up with MRA vs IA-DSA to assess whether in this setting MRA may replace IA-DSA.

**Methods**—We studied aneurysm occlusion on MRA in addition to follow-up IA-DSA in 310 patients with 341 coiled intracranial aneurysms. The observed sensitivity (82%) and specificity (89%) of MRA for detection of reopening with IA-DSA as a reference were used as input for a Markov decision-analytic model. Other determinants were derived from the literature. We compared life expectancy, quality-adjusted life-years (QALY), costs, and expected number of events for the two strategies.

**Results**—Follow-up with MRA yielded similar life expectancy (MRA, 26.66 years; IA-DSA, 26.63 years; difference, 0.03 years; 95% CI,  $-0.17$ – $0.23$ ) and QALY (MRA, 10.96; IA-DSA, 10.95; difference, 0.01 QALY; 95% CI,  $-0.05$ – $0.08$ ) at lower costs (MRA, \$7003; IA-DSA, \$8241 per patient; difference,  $-\$1238$ ; 95% CI,  $-2617$ – $36$ ). The expected number of events was comparable except for complications from IA-DSA.

**Conclusion**—MRA provided equivalent health benefits as IA-DSA and was cost-saving. MRA dominates and should replace routine IA-DSA to follow-up patients with coiled aneurysms. (*Stroke*. 2010;41:1736-1742.)

**Key Words:** cost-benefit analysis ■ digital subtraction angiography ■ intracranial aneurysm  
■ magnetic resonance angiography

Follow-up after occlusion of intracranial aneurysms with coils is required because reopening and subsequent rupture may occur.<sup>1–3</sup> Intra-arterial digital subtraction angiography (IA-DSA) is the standard modality to detect reopening after coiling but is invasive and irradiating.<sup>4</sup> Furthermore, the procedure may cause patient discomfort and requires substantial capacity of the angiography suite and inpatient clinic. Magnetic resonance angiography (MRA) is an alternative technique that is noninvasive, nonirradiating, and can be performed in an outpatient setting.

To investigate whether MRA can replace IA-DSA for follow-up of coiled patients, complete diagnostic evaluation of MRA is required. This should include assessment of its test characteristics, effect on clinical outcome, and cost-effectiveness.<sup>5</sup>

Although test characteristics have been reported, we could not find studies on effects on clinical outcome and cost-effectiveness in this clinical setting. In a large prospective series of patients, we have recently compared MRA and IA-DSA to assess reopening of coiled aneurysms.<sup>6</sup> This enabled us to use the observed test characteristics of MRA with IA-DSA as a reference to assess the expected changes in health benefits and costs incurred using MRA or IA-DSA.

## Materials and Methods

Using a cross-sectional design, we previously assessed the accuracy for detection of reopened aneurysms in MRA with IA-DSA as a reference in 310 coiled patients (mean age,  $51 \pm 12$ ; 71% women).<sup>6</sup> Unenhanced (time-of-flight) and contrast-enhanced MRA were performed in each patient in addition to routine IA-DSA. Two observers classified, independently from each other, the level of occlusion as

Received March 19, 2010; final revision received May 5, 2010; accepted May 8, 2010.

From the Department of Neurology (J.D.S., G.J.E.R.), Rudolf Magnus Institute of Neuroscience, University Medical Center, Utrecht, the Netherlands; the Department of Medical Technology Assessment (H.K.), Julius Center for Health Sciences and Primary Care, University Medical Center, Utrecht, the Netherlands; the Department of Epidemiology (E.B.), University Medical Center, Groningen, the Netherlands; the Department of Radiology (B.K.V.), Image Sciences Institute, University Medical Center, Utrecht, the Netherlands; the Department of Clinical Epidemiology (Y.G.), Julius Center for Health Sciences and Primary Care, University Medical Center, Utrecht, the Netherlands.


Correspondence to J.D. Schaafsma MD, Department of Neurology, Rudolf Magnus Institute of Neuroscience, University Medical Center, PO Box 85500, 3508 GA Utrecht, the Netherlands. E-mail j.d.schaafsma@umcutrecht.nl

© 2010 American Heart Association, Inc.

*Stroke* is available at <http://stroke.ahajournals.org>

DOI: 10.1161/STROKEAHA.110.585083





**Figure 2.** This simplified part of the decision tree used in the Markov model illustrates follow-up with MRA. TP, true-positive; FN, false-negative; TN, true-negative; FP, false-positive test result. The branch for IA-DSA is not displayed.

between the model parameters with associated uncertainty and changes in costs, effects in QALY, and cost-effectiveness.<sup>16</sup>

## Results

For the baseline model, life expectancy and QALY were in the same range for follow-up with MRA and for follow-up with IA-DSA, whereas MRA significantly reduced costs (Tables 2 and 3, Figure 3).

MRA induced health gain while saving costs in 67% of our samples and induced health gain while increasing costs in 1%. Conversely, MRA reduced health benefits at lower costs in 31% of our samples and reduced health benefits at increased costs in 1% (Figure 3).

Life expectancy was 39.42 years (95% CI, 39.03–39.80) after IA-DSA vs 39.47 years (95% CI, 39.04–39.91) after MRA for 35-year old patients, 26.63 years (95% CI, 26.31–26.95) after IA-DSA vs 26.66 years (95% CI, 26.35–26.98) after MRA for 50-year-old patients, and 15.81 years (95% CI, 15.59–16.02) after IA-DSA vs 15.82 years (95% CI, 15.59–16.02) after MRA for 65-year-old patients.

The number of events during follow-up with IA-DSA and MRA was not different except for case fatality and morbidity caused by IA-DSA (Table 2). The incidence of recurrent subarachnoid hemorrhage was not significantly higher for MRA than for IA-DSA. The difference in the overall case fatality and morbidity between the diagnostic strategies was 9 out of 5000 patients in favor of follow-up by MRA.

Scenario analyses for different ages, discount rates, and rupture rates of reopened aneurysms yielded similar results as for the baseline model. MRA remained cost-saving with a similar change in QALY compared to IA-DSA (Table 3). For 50-year-old patients MRA gained slightly more QALY in 68%, for 35-year-old patients in 72%, and for 65-year-old patients in 64% of the sampled cohorts. The probability of health gain by MRA increases with decreasing age. The cost-saving provided by MRA was similar in all our scenarios and was apparently not largely influenced by patient age, the rupture risk, or the discount rate used in the model.

Sensitivity analysis demonstrated that the distribution of input parameters did not significantly influence costs or QALY. We did not find an association between values of single-model parameters and cost-effectiveness estimates.

## Discussion

Follow-up after coiling of intracranial aneurysms by MRA results in similar health benefits but lower costs than follow-up by IA-DSA. The expected number of events was similar for both strategies, except for morbidity and case fatality caused by IA-DSA. Nevertheless, these complications did not have a major impact on cost-effectiveness because the total number of expected complications of IA-DSA remained small, particularly in comparison to patients with atherosclerosis.<sup>4</sup> The complication risk of MRA with contrast agent is extremely small.<sup>17,18</sup> We incorporated this small risk into the



**Table 1. Model Input Parameters**

Model Parameter	Value	95% CI/Range*	Distribution	Source
<b>Probabilities</b>				
Sensitivity MRA	0.82	0.72–0.89	Beta	CS
Specificity MRA	0.89	0.85–0.93	Beta	CS
<i>P</i> (case fatality of IA-DSA)	0.0006	0.0003–0.0010	Beta	4
<i>P</i> (morbidity of IA-DSA)	0.0010	0.0006–0.0015	Beta	4
<i>P</i> (case fatality of MRA)	0.0000000	...	...	17, 18
<i>P</i> (morbidity of MRA)	0.0000053	0.0000026–0.0000088	Beta	17, 18
<i>P</i> (case fatality recoiling)	0.0056	0.0018–0.0120	Beta	24–27
<i>P</i> (morbidity recoiling)	0.011	0.004–0.021	Beta	26, 27
<i>P</i> (case fatality rebleeding)				28
30–39 y	0.182	0.162–0.204	Beta	
40–49 y	0.225	0.211–0.240	Beta	
50–59 y	0.249	0.236–0.262	Beta	
60–69 y	0.317	0.310–0.324	Beta	
70–79 y	0.455	0.438–0.472	Beta	
>80 y	0.576	0.553–0.598	Beta	
<i>P</i> (morbidity rebleeding)	0.09	0.08–0.11	Beta	29, 30
<i>P</i> (case fatality disabled patients) per year	0.24	0.10–0.36	Uniform	28, 31
<i>P</i> (reopening) up to 6 mo after coiling†	0.119	0.097–0.144	Beta	10, 11, 13
<i>P</i> (reopening) 6–18 mo after coiling†	0.055	0.019–0.110	Beta	9, 11, 14
<i>P</i> (reopening) 18 mo–6 y after coiling†	0.036	0.010–0.078	Beta	12
<i>P</i> (recoiling reopened aneurysm)	0.58	0.47–0.68	Beta	CS
<i>P</i> (rupture risk reopening) per year	0.017	0.014–0.020	Uniform	10, 12, 22, 23, 32
<b>Costs (\$)</b>				
Costs IA-DSA	\$838	...	...	33
Costs MRA	\$371	...	...	33
Costs of rebleeding including recoiling	\$36 920	...	...	34, 35
Costs of elective recoiling	\$12 646	...	...	33
Costs nursing home per year	\$107 711	...	...	15
Costs of patient death	\$3585	...	...	36
<b>Utilities</b>				
Well after subarachnoid hemorrhage	0.72	0.65–0.80	Triangular	37
Disabled	0.25	0.21–0.30	Triangular	37
Dead	0	...	...	
<b>Discounting</b>				
Cost discount per year	4%	...	...	...
Effect discount per year	4%	...	...	...

CS indicates data obtained from our clinical study; IA-DSA, intra-arterial digital subtraction angiography; MRA, magnetic resonance imaging.

\*95% CI for beta distribution; range for uniform/triangular distribution.

†A mathematical function based on these results was used (provided online). All rates obtained were converted to probabilities per month.

model, even though in our clinical study and other studies contrast-enhanced MRA did not provide significant additional information for unenhanced MRA.<sup>19,20</sup> So, the administration of contrast agent is often unnecessary, which decreases the morbidity risk of MRA even further. Moreover, those reopened aneurysms on IA-DSA that were not identified on MRA did not significantly increase the expected incidence of subarachnoid hemorrhage for patients

followed-up with MRA. As a result of less than optimal quality of life after subarachnoid hemorrhage, life expectancy considerably surpasses the number of QALY, regardless of the strategy.

MRA appeared to be cost-saving. Because the number of events does not largely differ between the two strategies, the difference in costs is likely to be caused by the lower costs of MRA compared to the IA-DSA procedure. When cost reduc-

**Table 2. Costs and Effects for Follow-Up With IA-DSA vs MRA**

	IA-DSA	MRA	Difference	95% CI
Life expectancy per patient, y	26.63	26.66	0.03	−0.17–0.23
QALY per patient	10.95	10.96	0.01	−0.05–0.08
Total case fatality strategy, n	125	120	−5	−24–13
Total morbidity strategy, n	38	34	−4	−16–11
Case fatality recoiling, n	5	5	0	−3–2
Morbidity recoiling, n	9	9	0	−4–3
Case fatality test, n	7	0	−7	−14–−2*
Morbidity test, n	11	0	−11	−20–−4*
Reopened aneurysms, n	1359	1360	1	−28–31
Recoiling procedures, n	728	718	−10	−29–7
Rebleedings, n	191	196	5	−12–22
Costs per patient	\$8241	\$7003	\$−1238	−2617–−36*

The cost-effectiveness estimates based on simulations of 2500 cohorts of 5000 patients each.

IA-DSA, intra-arterial digital subtraction angiography; MRA, magnetic resonance imaging; QALY, quality-adjusted life-year.

\*Statistically significant.

tion is not taken into consideration, there is still a substantial chance that MRA is the preferred strategy with a small gain in QALY, especially for younger patients.

We could not find other studies on cost-effectiveness of MRA vs IA-DSA after coiling. Using a Markov model, we integrated the best available evidence for computation of the expected long-term outcomes. A diagnostic, randomized, clinical trial, although theoretically more accurate, would be infeasible because a large number of participants and long follow-up is required to ascertain the incidence of rupture of undetected or untreated reopened aneurysms.<sup>5</sup>

We intended to construct a detailed Markov model that appropriately reflects clinical practice, although we faced some limitations. First, IA-DSA is not a perfect reference standard for follow-up of coiled aneurysms. For example, the 2-dimensional images restrict visualization of residual flow in case of superimposition of coils or surrounding arteries.<sup>21</sup> Consequently, discrepant results on MRA had to be labeled as “false-positive” or “false-negative,” whereas MRA may provide the more realistic visualization. Thus, the model represented the least favorable scenario for MRA and therefore may underestimate its diagnostic performance. MRA still appeared dominant, thus strengthening the conclusion that MRA may replace IA-DSA.

Second, input parameters originated from our clinical study and from the literature. Because coiling has been available since 1992, only limited data on reopening and subsequent rupture rates more than 5 to 10 years after coiling are available.<sup>1,2,10,22,23</sup> Reopening rates could only be estimated from a few studies with a systematic long-term follow-up at fixed time intervals.<sup>3,9–14</sup>

Third, for the model, we assumed a similar rupture rate for aneurysms that reopened after follow-up for untreatable and for undetected reopened aneurysms, whereas the actual rupture risks may differ in each situation. Undetected reopened aneurysms in our clinical study were smaller and therefore probably had a lower rupture rate than larger reopened aneurysms that are left untreated. By assuming a similar rupture rate, we overestimated the health loss from undetected aneurysms when using MRA and therefore underestimated the health benefits provided by MRA. Repeated analyses for different rupture rates resulted in marginal and similar changes in QALY and costs. So, the uncertainty around the exact rupture rate did not influence the cost-effectiveness of MRA compared to IA-DSA. We furthermore assumed that reopened aneurysms never occlude spontaneously. In case of spontaneous occlusion, the potential hazard of a nonidentified reopened aneurysm on MRA would be smaller. So, again, we applied the least favorable scenario for MRA to avoid positive bias.


**Table 3. Scenario Analyses**

Age, y	DR, %	Rupture Risk*	Costs, \$		QALY		ΔCosts	95% CI	ΔQALY	95% CI
			IA-DSA	MRA	IA-DSA	MRA				
35	4	0.017	8240	7001	13.40	13.42	−1238	(−2663–−75)	0.02	(−0.04–0.08)
50	4	0.017	8241	7003	10.95	10.96	−1238	(−2617–−36)	0.01	(−0.05–0.08)
65	4	0.017	8290	7015	7.95	7.96	−1275	(−2523–−199)	0.01	(−0.04–0.06)
35	1.5	0.017	11 322	10 068	20.76	20.79	−1254	(−2899–290)	0.03	(−0.09–0.13)
50	1.5	0.017	10 438	9153	15.29	15.31	−1284	(−2979–72)	0.02	(−0.08–0.12)
65	1.5	0.017	9743	8403	9.90	9.91	−1340	(−2894–−50)	0.01	(−0.06–0.09)
50	4	0.034	9284	8056	10.90	10.91	−1227	(−2588–−6)	0.01	(−0.05–0.07)
50	4	0.014	7897	6653	10.96	10.98	−1244	(−2295–−108)	0.02	(−0.05–0.09)
50	4	0.009	7546	6291	10.98	10.99	−1255	(−2492–−61)	0.01	(−0.05–0.07)
50	4	0.005	7135	5871	11.00	11.01	−1264	(−2677–−122)	0.01	(−0.05–0.08)

Each analysis comprised 1000 simulations for 5000 patients.

DR, discount rate; IA-DSA, intra-arterial digital subtraction angiography; MRA, magnetic resonance imaging; QALY, quality-adjusted life-year.

\*Rupture risk of reopened aneurysm per year.



**Figure 3.** This cost-effectiveness graph shows the increment in QALY vs the increment in costs induced by MRA for 2500 simulations. Each dot represents one simulation for 5000 patients for which input values were sampled from the parameter distributions given in Table 1. The percentages of samples per quadrant are indicated.

Fourth, we did not evaluate the influence of uncertainty in costs on the cost-effectiveness of MRA vs IA-DSA because insufficient evidence was available to define their distribution. Finally, because information on actual dependencies between model parameters could not be obtained, all parameters were, by necessity, assumed to be independent in our probabilistic sensitivity analysis. This assumption may not hold for all parameters, eg, for sensitivity and specificity. Nevertheless, because the sensitivity analyses showed overall robust outcomes, we feel that our general conclusion remains justified.

MRA is a safe technique that can be performed in an outpatient setting. Our results show that the consequences of misdiagnosis by MRA outweigh the complications caused by IA-DSA and that MRA reduces costs. We therefore recommend using MRA instead of IA-DSA to follow-up coiled patients. The exact timing of reopening and subsequent rupture after coiling is unclear. Additional studies on timing of follow-up MRA are warranted to assess the short-term and long-term evolution of coiled aneurysms.

### Conclusion

Cost-effectiveness analysis by Markov modeling shows that potential consequences of misdiagnosis by MRA will be offset by the direct risk of complications associated with IA-DSA, and MRA will reduce costs considerably. Patients therefore should be followed-up by MRA instead of IA-DSA to detect reopening after coiling of intracranial aneurysms.

### Sources of Funding

J.D.S. was supported by a grant from the Netherlands Organization for Scientific Research (NWO) and the Netherlands Organization for Health Research and Development (ZonMw; grant 945-04-310).

### Disclosures

None.

### References

1. Molyneux AJ, Kerr RS, Birks J, Ramzi N, Yarnold J, Sneade M, Rischmiller J. Risk of recurrent subarachnoid haemorrhage, death, or dependence and standardised mortality ratios after clipping or coiling of an intracranial aneurysm in the International Subarachnoid Aneurysm Trial (ISAT): long-term follow-up. *Lancet Neurol*. 2009;8:427–433.
2. CARAT investigators. Rates of delayed rebleeding from intracranial aneurysms are low after surgical and endovascular treatment. *Stroke*. 2006;37:1437–1442.
3. Ferns SP, Sprengers ME, van Rooij WJ, Rinkel GJ, van Rijn JC, Bipat S, Sluzewski M, Majoie CB. Coiling of intracranial aneurysms: a systematic review on initial occlusion and reopening and retreatment rates. *Stroke*. 2009;40:e523–e529.
4. Kaufmann TJ, Huston J III, Mandrekar JN, Schleck CD, Thielen KR, Kallmes DF. Complications of diagnostic cerebral angiography: evaluation of 19,826 consecutive patients. *Radiology*. 2007;243:812–819.
5. Schaafsma JD, van der Graaf Y, Rinkel GJ, Buskens E. Decision analysis to complete diagnostic research by closing the gap between test characteristics and cost-effectiveness. *J Clin Epidemiol*. 2009;62:1248–1252.
6. Schaafsma JD, Velthuis BK, Majoie CBLM, van den Berg R, Brouwer PA, Barkhof F, Eshghi O, De Kort GAP, Lo TH, Witkamp TD, Sprengers MES, Van Walderveen MA, Bot JC, Sanchez E, Vandertop P, van Gijn J, Buskens E, van der Graaf Y, Rinkel GJE. Test characteristics of magnetic resonance angiography in the follow-up of patients with coiled intracranial aneurysms: a cross-sectional multi-center study. *Radiology*. 2010.
7. Briggs A, Sculpher M. An introduction to Markov modelling for economic evaluation. *Pharmacoeconomics*. 1998;13:397–409.
8. Wermer MJ, Greebe P, Algra A, Rinkel GJ. Long-term mortality and vascular event risk after aneurysmal subarachnoid haemorrhage. *J Neurol Neurosurg Psychiatry*. 2009;80:1399–1401.
9. Ries T, Siemonsen S, Thomalla G, Grzyska U, Zeumer H, Fiehler J. Long-term follow-up of cerebral aneurysms after endovascular therapy prediction and outcome of retreatment. *AJNR Am J Neuroradiol*. 2007;28:1755–1761.
10. Sluzewski M, van Rooij WJ, Rinkel GJ, Wijnalda D. Endovascular treatment of ruptured intracranial aneurysms with detachable coils: long-term clinical and serial angiographic results. *Radiology*. 2003;227:720–724.
11. Raymond J, Guilbert F, Weill A, Georganos SA, Juravsky L, Lambert A, Lamoureaux J, Chagnon M, Roy D. Long-term angiographic recurrences after selective endovascular treatment of aneurysms with detachable coils. *Stroke*. 2003;34:1398–1403.
12. Sprengers ME, Schaafsma J, van Rooij WJ, Sluzewski M, Rinkel GJ, Velthuis BK, van Rijn JC, Majoie CB. Stability of intracranial aneurysms



- adequately occluded 6 months after coiling: a 3T MR angiography multicenter long-term follow-up study. *AJNR Am J Neuroradiol*. 2008;29:1768–1774.
13. Kang HS, Han MH, Kwon BJ, Kwon OK, Kim SH. Repeat endovascular treatment in post-embolization recurrent intracranial aneurysms. *Neurosurgery*. 2006;58:60–70.
  14. Murayama Y, Nien YL, Duckwiler G, Gobin YP, Jahan R, Frazee J, Martin N, Vinuela F. Guglielmi detachable coil embolization of cerebral aneurysms: 11 years' experience. *J Neurosurg*. 2003;98:959–966.
  15. Oostenbrink J, Bouwmans C, Koopmanschap M, Rutten F. Instructions for cost assessment, methods and directive of prices for economic evaluation in health care (in Dutch). Amstelveen, The Netherlands: Health Care Insurance Board; 2004.
  16. Ades AE, Claxton K, Sculpher M. Evidence synthesis, parameter correlation and probabilistic sensitivity analysis. *Health Econ*. 2006;15:373–381.
  17. Murphy KP, Szopinski KT, Cohan RH, Mermillod B, Ellis JH. Occurrence of adverse reactions to gadolinium-based contrast material and management of patients at increased risk: a survey of the American Society of Neuroradiology Fellowship Directors. *Acad Radiol*. 1999;6:656–664.
  18. Niendorf HP, Dinger JC, Haustein J, Cornelius I, Alhassan A, Clauss W. Tolerance data of Gd-DTPA: a review. *Eur J Radiol*. 1991;13:15–20.
  19. Kwee TC, Kwee RM. MR angiography in the follow-up of intracranial aneurysms treated with Guglielmi detachable coils: systematic review and meta-analysis. *Neuroradiology*. 2007;49:703–713.
  20. Sprengers ME, Schaafsma JD, van Rooij WJ, van den Berg R, Rinkel GJ, Akkerman EM, Ferns SP, Majoie CB. Evaluation of the occlusion status of coiled intracranial aneurysms with MR angiography at 3T: is contrast enhancement necessary? *AJNR Am J Neuroradiol*. 2009;30:1665–1671.
  21. Agid R, Willinsky RA, Lee SK, Terbrugge KG, Farb RI. Characterization of aneurysm remnants after endovascular treatment: contrast-enhanced MR angiography vs catheter digital subtraction angiography. *AJNR Am J Neuroradiol*. 2008;29:1570–1574.
  22. Byrne JV, Sohn MJ, Molyneux AJ, Chir B. Five-year experience in using coil embolization for ruptured intracranial aneurysms: outcomes and incidence of late rebleeding. *J Neurosurg*. 1999;90:656–663.
  23. Schaafsma JD, Sprengers ME, van Rooij WJ, Sluzewski M, Majoie CB, Wermer MJ, Rinkel GJ. Long-term recurrent subarachnoid hemorrhage after adequate coiling vs clipping of ruptured intracranial aneurysms. *Stroke*. 2009;40:1758–1763.
  24. Gonzalez N, Murayama Y, Nien YL, Martin N, Frazee J, Duckwiler G, Jahan R, Gobin YP, Vinuela F. Treatment of unruptured aneurysms with GDCs: clinical experience with 247 aneurysms. *AJNR Am J Neuroradiol*. 2004;25:577–583.
  25. Grunwald IQ, Papanagiotou P, Politi M, Struffert T, Roth C, Reith W. Endovascular treatment of unruptured intracranial aneurysms: occurrence of thromboembolic events. *Neurosurgery*. 2006;58:612–618.
  26. Lanterna LA, Tredici G, Dimitrov BD, Biroli F. Treatment of unruptured cerebral aneurysms by embolization with guglielmi detachable coils: case-fatality, morbidity, and effectiveness in preventing bleeding—a systematic review of the literature. *Neurosurgery*. 2004;55:767–775.
  27. van Rooij WJ, Sluzewski M. Procedural morbidity and mortality of elective coil treatment of unruptured intracranial aneurysms. *AJNR Am J Neuroradiol*. 2006;27:1678–1680.
  28. Koffijberg H, Buskens E, Granath F, Adami J, Ekblom A, Rinkel GJ, Blomqvist P. Subarachnoid haemorrhage in Sweden 1987–2002: regional incidence and case fatality rates. *J Neurol Neurosurg Psychiatry*. 2008;79:294–299.
  29. Hop JW, Rinkel GJ, Algra A, van Gijn J. Case-fatality rates and functional outcome after subarachnoid hemorrhage: a systematic review. *Stroke*. 1997;28:660–664.
  30. Ronkainen A, Niskanen M, Piironen R, Hernesniemi J. Familial subarachnoid hemorrhage. Outcome study. *Stroke*. 1999;30:1099–1102.
  31. van Dijk PT, Mehr DR, Ooms ME, Madsen R, Petroski G, Frijters DH, Pot AM, Ribbe MW. Comorbidity and 1-year mortality risks in nursing home residents. *J Am Geriatr Soc*. 2005;53:660–665.
  32. Johnston SC, Dowd CF, Higashida RT, Lawton MT, Duckwiler GR, Gress DR. Predictors of rehemorrhage after treatment of ruptured intracranial aneurysms: the Cerebral Aneurysm Rerupture After Treatment (CARAT) study. *Stroke*. 2008;39:120–125.
  33. Halkes PH, Wermer MJ, Rinkel GJ, Buskens E. Direct costs of surgical clipping and endovascular coiling of unruptured intracranial aneurysms. *Cerebrovasc Dis*. 2006;22:40–45.
  34. Roos YB, Dijkgraaf MG, Albrecht KW, Beenen LF, Groen RJ, de Haan RJ, Vermeulen M. Direct costs of modern treatment of aneurysmal subarachnoid hemorrhage in the first year after diagnosis. *Stroke*. 2002;33:1595–1599.
  35. Wolstenholme J, Rivero-Arias O, Gray A, Molyneux AJ, Kerr RS, Yarnold JA, Sneade M. Treatment pathways, resource use, and costs of endovascular coiling vs surgical clipping after aSAH. *Stroke*. 2008;39:111–119.
  36. Buskens E, Nederkoorn PJ, Buijs-Van der Woude T, Mali WP, Kappelle LJ, Eikelboom BC, van der Graaf Y, Hunink MG. Imaging of carotid arteries in symptomatic patients: cost-effectiveness of diagnostic strategies. *Radiology*. 2004;233:101–112.
  37. Post PN, Stiggelbout AM, Wakker PP. The utility of health states after stroke: a systematic review of the literature. *Stroke*. 2001;32:1425–1429.